

INSECT SCIENCE AND ITS APPLICATION

The International Journal of Tropical Insect Science

Editor-in-Chief

Thomas R. Odhiambo

The International Centre of Insect Physiology and Ecology (ICIPE)

List of Contents, Author and Subject
Index for Volume 12, 1991

CAB INTERNATIONAL
Library Services Centre

- 7 JUL 1992

ICIPE SCIENCE PRESS

Insect Science and its Application

The International Journal of Tropical Insect Science

Sponsored by the International Centre of Insect Physiology and Ecology (ICIPE), and the African Association of Insect Scientists (AAIS)

Editor-in-Chief: Thomas R. Odhiambo

Associate Editor: K. N. Saxena

Senior Scientific Editor: Serah W. Mwanycky

The International Centre of Insect Physiology and Ecology (ICIPE), P.O. Box 30772, Nairobi, Kenya

Editorial Advisory Board

F. T. Abushama, Kuwait, State of Kuwait

H. C. Agarwal, Delhi, India

J. Allen, Saskatoon, Canada

J. K. O. Ampofo, Arusha, Tanzania

A. Attygalle, Ithaca, New York, USA

J. L. Auclair, Montreal, Canada

R. Galun, Jerusalem, Israel

I. Fagoonee, Reduit, Mauritius

K. M. Harris, London, UK

T. Hidaka, Kyoto, Japan

H. Hirumi, Nairobi, Kenya

A. G. Ibrahim, Serdang, Selangor, Malaysia

R. Kumar, Port Harcourt, Nigeria

V. M. Labeyrie, Paris, France

A. Mansingh, Kingston, Jamaica

F. G. Maxwell, College Station, Texas, USA

A. I. Mohyuddin, Rawalpindi, Pakistan

J. Mouchet, Paris, France

R. W. Mwangi, Nairobi, Kenya

L. M. Schoonhoven, Wageningen, The Netherlands

P. Singh, Auckland, New Zealand

K. Slama, Praha, Czechoslovakia

G. L. Teetes, College Station, Texas, USA

S. A. Toye, Ibadan, Nigeria

H. F. van Emden, Reading, UK

A. van Huis, Wageningen, The Netherlands

S. Yagi, Tsukuba, Ibaraki, Japan

PUBLISHED BIMONTHLY

Annual Subscription

For libraries, university departments, government laboratories, industrial and other multiple reader institutions US\$ 210.00 (1992); 2-year rate (1992–1993) US\$ 410.00 (including postage and insurance), private individuals US\$ 80.00 (1992); 2-year rate US\$ 150.00 (1992–1993). *Specially Reduced Rates for Individuals:* In the interest of maximizing the dissemination of the research results published in this important international journal we have established a two-tier price structure. Any individual whose institution takes out a library subscription may purchase a second or additional subscriptions for personal use at a much reduced rate of US\$ 70.00 (1992); 2-year rate (1992–1993) US\$ 140.00. Members of the International Centre of Insect Physiology and Ecology (ICIPE) and the African Association of Insect Scientists (AAIS) may purchase a subscription at the special rate of US\$ 20.00 (1992); 2-year rate (1992–1993) US\$ 40.00.

Application forms for membership of the AAIS (registration fee US\$ 20.00; annual membership fee US\$ 20.00) may be obtained from: Hon. Secretary, African Association of Insect Scientists, P.O. Box 59862, Nairobi, Kenya.

Prices are subject to amendment without notice

Copyright © 1992—ICIPE Science Press, The International Centre of Insect Physiology and Ecology

It is a condition of publication that manuscripts submitted to this journal have not been published and will not be simultaneously submitted or published elsewhere. By submitting a manuscript, the authors agree that the copyright for their article is transferred to the publisher if and when the article is accepted for publication. However, assignment of copyright is not required from authors who work for organizations which do not permit such assignment. The copyright covers the exclusive rights to reproduce and distribute the article, including reprints, photographic reproductions, microform or any other reproductions of similar nature and translations. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, electrostatic, magnetic tape, mechanical photocopying, recording or otherwise, without permission in writing from the copyright holder.

Photocopying Information for users in the USA

The Item-fee Code for this publication indicates that authorization to photocopy items for internal or personal use is granted by the copyright holder for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided the stated fee for copying beyond that permitted by Section 107 or 108 of the US Copyright Law, is paid. The appropriate remittance of \$3.00 per copy per article is paid directly to the Copyright Clearance Center Inc., 27 Congress Street, Salem, MA 01970.

Permission for other use

The copyright owner's consent does not extend to copying for general distribution, for promotion, for creating new works or for resale. Specific written permission must be obtained from the publisher of such copying.

The item-fee for this publication is 0191-9040/92 \$3.00+0.00

ICIPE SCIENCE PRESS
P. O. Box 72913, Nairobi, Kenya, East Africa

CONTENTS OF VOLUME 12

VOLUME 12 NUMBER 1/2/3

SPECIAL ISSUE

CURRENT TOPICS IN TROPICAL ENTOMOLOGY

T. R. ODHIAMBO	1	Foreword
D. J. GREATHEAD	3	Biological control in the tropics: Present opportunities and future prospects
V. K. GUPTA	9	The parasitic Hymenoptera and biological control and our knowledge of the African Ichneumonidae
A. I. MOHYUDDIN	19	Utilization of natural enemies for the control of insect pests of sugar-cane
S. T. MURPHY and D. K. RANGI	27	The use of African wasp, <i>Proropes nasuta</i> for the control of the coffee berry borer, <i>Hypothenemus hampei</i> in Mexico and Ecuador: The introduction programme
J. B. OKEYO-OWUOR, G. W. OLOO and P. O. AGWARO	35	Natural enemies of the legume pod borer <i>Maruca testulalis</i> Geyer (Lepidoptera: Pyralidae) in small scale farming systems of Western Kenya
E.-D. N. UMEH	43	Natural enemies in the control of pests of cassava <i>Manihot esculenta</i> Crantz
M. O. ODINDO	51	Management of cereal stem borers, especially <i>Chilo partellus</i> , using microsporidia
M. BROWNBIDGE	57	Native <i>Bacillus thuringiensis</i> isolates for the management of Lepidopteran cereal pests
N. K. MANIANIA	63	Potential of some fungal pathogens for the control of pests in the tropics
W. S. BOWERS and F. ORTEGO	71	Evaluation of juvenoid insect growth regulators on <i>Schistocerca americana</i>
R. KUMAR and N. O. OKONRONKWO	77	Effectiveness of plant oils against some Bostrychidae infesting cereals in storage
A. VAN HUIS	87	Biological methods of bruchid control in the tropics: A review
L. E. N. JACKAI and I. O. OYEDIRAN	103	The potential of neem <i>Azadirachta indica</i> A. Juss. for controlling post-flowering pests of cowpea <i>Vigna unguiculata</i> Walp. — I. The pod borer, <i>Maruca testulalis</i>
I. JAVAID	111	Application of insecticides on cotton in Zambia: Comparison of application techniques
I. JAVAID	127	Application of insecticides on cotton in Zambia: Electrodyn swath rows

R. G. PEGRAM, A. D. JAMES, G. P. M. OOSTERWIJK and H. G. B. CHIZYUKA	139	The economic impact of cattle tick control in Central Africa
R. S. PATHAK	147	Genetic expression of the spotted stem borer, <i>Chilo partellus</i> (Swinhoe) resistance in three maize crosses
F. M. E. WANJALA and M. W. KINUTHIA	153	Relating incidence of <i>Icerya pattersoni</i> Newst (Homoptera: Margorodidae) to characteristics of veins of coffee leaves
D. P. GIGA and U. M. MAZARURA	159	Levels of resistance to the maize weevil, <i>Sitophilus zeamais</i> (Motsch.) in exotic, local open-pollinated and hybrid maize germplasm
S. OGHIAKHE, L. E. N. JACKAI and W. A. MAKANJUOLA	171	Anatomical parameters of cowpea, <i>Vigna unguiculata</i> (L.) Walp. stem and pod wall resistance to the legume pod borer <i>Maruca testulalis</i> Geyer (Lepidoptera: Pyralidae)
C. S. GOLD and J. A. WIGHTMAN	177	Effects of intercropping groundnut with sunnhemp on termite incidence and damage in India
K. F. NWANZE, Y. V. R. REDDY, S. L. TANEJA, H. C. SHARMA and B. L. AGRAWAL	183	Evaluating sorghum genotypes for multiple insect resistance
F. P. NEUPANE	189	Effect of soybean hosts on the development of the soybean hairy caterpillar, <i>Spilarctia casignata</i> Koll. (Lepidoptera: Arctiidae)
S. OGHIAKHE, L. E. N. JACKAI and W. A. MAKANJUOLA	193	Cowpea plant architecture in relation to infestation and damage by the legume pod borer <i>Maruca testulalis</i> Geyer (Lepidoptera: Pyralidae) — I. Effect of canopy structure and pod position
L. G. PHILIPPE	201	Niche breadth and feeding in tropical grasshoppers
P. MOYAL and M. TRAN	209	Cob borer <i>Mussidia nigrivenella</i> (Lepidoptera: Pyralidae) of maize in Ivory Coast. I — Morphological and biological data
P. MOYAL and M. TRAN	215	Cob borer <i>Mussidia nigrivenella</i> Ragonot (Lepidoptera: Pyralidae) of maize in Ivory Coast. II — Ecological data
D. WOOL, D. GERLING, A. BELLOTTI, F. MORALES and B. NOLT	225	Spatial and temporal genetic variation in populations of the whitefly <i>Bemisia tabaci</i> (Genn.) in Israel and Columbia: An interim report
B. T. NYAMBO	231	The pest status of <i>Zonocerus elegans</i> (Thunberg) (Orthoptera: Acridoidea) in Kilosa District in Tanzania with some suggestions on control strategies
Z. T. DABROWSKI and P. S. CWIKLA	237	Some new observations on <i>Cicadulina</i> leafhoppers taxonomic structures
Z. T. DABROWSKI, F. NWILENE and R. KUMAR	249	First regular observations on leafhoppers, <i>Cicadulina</i> spp., vectors of maize streak virus (MSV) in southeastern Nigeria
Y. ZHENG-XIANG and L. RI-ZHI	263	A study on the occurrence tendency of <i>Tryporyza incertulas</i> (Walker) by grey prediction
K. V. SESHU REDDY and K. O. S. SUM	269	Determination of economic injury level of the stem borer, <i>Chilo partellus</i> (Swinhoe) in maize, <i>Zea mays</i> L.
G. F. RAJENDRAM	275	Electrophoretic study of enzymes from <i>Sogatella furcifera</i> Horvath population from Sri Lanka

A. M. TH. BEENAKKERS	279	Adipokinetic hormones and lipoprotein inter-conversions during locust flight
W. TAKKEN	287	The role of olfaction in host-seeking of mosquitoes: A review
C. S. GOLD, J. A. WIGHTMAN and M. P. PIMBERT	297	Effects of mulches on foraging behaviour of <i>Microtermes obesi</i> and <i>Odontotermes</i> spp. in India
S. B. SAGNIA	305	Training in entomology for middle-level technicians at the CILSS regional crop protection training centre in Niger
K. M. HARRIS	311	Plenary Lecture: Summing up
Instructions to Authors	i	
Author Checklist	iv	

VOLUME 12 NUMBER 4

R. T. GAHUKAR	313	<i>Mini Review Article</i> Control of cotton insect and mite pests in sub-tropical Africa: Current status and future needs
C. PETER and B. V. DAVID	339	<i>Research Articles</i> Biology of <i>Goniozus sensorius</i> Gordh (Hymenoptera: Bethyridae) a parasitoid of the pumpkin caterpillar, <i>Diaphania indica</i> (Saunders) (Lepidoptera: Pyralidae)
J. SRIKANTH and N. A. PUSHPALATHA	347	Status of biological control of <i>Parthenium hysterophorus</i> L. in India: A review
M. D. AWOYEMI	361	Preliminary observations on the incidence of insect pests and related damage to stored dried fish in Kainji Lake area
S. R. TRIPATHI and R. SINGH	367	Population dynamics of <i>Helicoverpa armigera</i> (Hubner) (Lepidoptera: Noctuidae)
M. LULU, W. NIGATU, T. GEZAHEGN and D. TILAHUN	375	Inversion polymorphisms in <i>Anopheles arabiensis</i> (Patton) in five selected localities from east, south and southwest Ethiopia
S. T. NESBITT, B. C. NJAU and L. H. OTIENO	379	Epizootiology of trypanosomiasis in Lambwe Valley, Kenya, East Africa
N. BHATT and R. SINGH	385	Bionomics of an aphidiid parasitoid, <i>Trioxys indicus</i> Subba Rao and Sharma. 35. Influence of food plants on the life table statistics of the parasitoid through its host <i>Aphis gossypii</i> Glover
K. P. SALIN, C. KAILASAM and M. GOPALAN	391	A mathematical model for population growth of buprofezin treated brown planthopper <i>Nilaparvata lugens</i> Stal. (Homoptera: Delphacidae)
O. O. DIPEOLU and V. KHAN	395	Studies on the dissemination of coccidia oocysts by <i>Musca</i> Linne, 1758 species
K. J. MBATA	401	Plant associations and rates of population interchange between two food plant areas of <i>Acanthoplus speiseri</i> Brancsik (Orthoptera: Tettigoniidae, Heterodinae)
K. J. MBATA	409	Spatial dispersion patterns of the armoured ground cricket, <i>Acanthoplus speiseri</i> Brancsik at the University of Zambia campus (Orthoptera: Tettigoniidae, Heterodinae)

S. P. SINGH	419	Biometrical observations on sorghum stem borer, <i>Chilo partellus</i> (Swinhoe) under fluctuating and constant temperature conditions
J. B. OKEYO-OWUOR and G. W. OLOO	423	Life tables, key factor analysis and density relations in natural populations of the legume pod borer <i>Maruca testulalis</i> Geyer (Lepidoptera: Pyralidae) in Western Kenya
K. C. NARAYANASWAMY, T. K. NARAYANASWAMY, M. C. DEVAIAH, B. L. VISWESWARA GOWDA and R. GOVINDAN	433	Evaluation of bivoltine single and double cross hybrids of silkworm, <i>Bombyx mori</i> L. for grainage parameters
G. V. PRASAD REDDY and C. DEVARAJ URS	439	Growth regulator activity of the xerophytic perennial plant, <i>Agave cantala</i> Roxb. on diamondback moth, <i>Plutella xylostella</i> (L.) (Lepidoptera: Yponomeutidae)
G. V. RANGA RAO, J. A. WIGHTMAN and D. V. RANGA RAO	443	Monitoring <i>Spodoptera litura</i> (F) (Lepidoptera: Noctuidae) using sex attractant traps: Effect of trap height and time of the night on moth catch
G. O. OLATUNDE, J. A. ODEBIYI and L. E. N. JACKAI	449	Cowpea antixenosis to the pod sucking bug, <i>Clavigralla tomentosicollis</i> Stal. (Hemiptera: Coreidae)
G. O. OLATUNDE, J. A. ODEBIYI, H. S. CHIANG and L. E. N. JACKAI	455	Identification of sources of resistance in cowpea, <i>Vigna unguiculata</i> L. Walp. to <i>Clavigralla tomentosicollis</i> Stal. (Hemiptera: Coreidae)
J. H. NDERITU, H. Y. KAYUMBO and J. M. MUEKE	463	Evaluation of bean cultivars for resistance to beanflies in single plantings at different seasons
R. H. S. RAJAPAKSE, T. R. ASHLEY and van H. WADDILL	473	Interspecific competition between parasitoids of the fall armyworm, <i>Spodoptera frugiperda</i> (J. E. Smith) (Lepidoptera: Noctuidae)
S. A. A. AMER, N. Z. DIMETRY and A. S. REDA	481	Toxicity of green marine algae to the two-spotted spider mite <i>Tetranychus urticae</i> Koch
D. SINGH and A. K. SINGH	487	Repellent and insecticidal properties of essential oils against housefly, <i>Musca domestica</i> L.
Book Reviews	493	
Announcement	500	Symposium
Instructions to Authors	i	
Author Checklist	iv	

VOLUME 12 NUMBER 5/6

SPECIAL ISSUE

T. R. ODHIAMBO	501	Foreword
----------------	-----	----------

PART I

SYMPOSIUM ON INTEGRATED PEST MANAGEMENT AND ENVIRONMENTAL CONSERVATION

ECOLOGY, BIOLOGY AND BIONOMICS

D. SECK	507	Etude de l'infestation initiale de <i>Sitotroga cerealella</i> Oliv. (Lepidoptera, Gelechiidae) en fonction de la localisation des champs de mil, <i>Pennisetum typhoides</i> (L) Lecke
M. J. N. OKWAKOL	511	Fauna associated with <i>Cubitermes testaceus</i> mounds (Isoptera: Termitidae)

INSECT PHYSIOLOGY

- J. KONGORO, T. R. ODHAMBO and J. R. MAINOYA 517 An ultrastructural study of Malpighian tubules of the stable fly, *Stomoxys calcitrans* (L.) (Diptera: Muscidae): Cell types and structural zonation

PEST MANAGEMENT STRATEGIES AND PRACTICES

- M. O. ODINDO 529 Pest and vector management in Africa
- E. M. TUKAHIRWA 535 Integrated pest management: Options for Uganda
- OLE ZETHNER 541 Integrated pest and land management experiences and perspectives in The Gambia
- EL TIGANI M. EL AMIN and M. A. AHMED 547 Strategies for integrated cotton pest control in the Sudan. 1 — Cultural and legislative measures
- R. S. PATHAK 553 Plant genetics in pest management
- D. DAKOUO, S. NACRO et B. BACYE 565 Mise au point d'un systeme de lutte rationnelle contre les insectes ravageurs sur les perimetres rizicoles irrigues au Burkina Faso
- R. J. LAVIGNE 571 Stored grain insects in underground storage pits in Somalia and their control
- C. W. BALIDDAWA 579 Agricultural extension and pest control: A case study of Iganga District
- M. I. EZUEH 585 Prospects for cultural and biological control of cowpea pests
- B. S. KILONZO 593 The effects of permethrin on *Ctenocephalides felis*, the commonest flea pest of livestock in Tanzania
- B. S. TOGUEBAYE 599 Effet du parasitisme par *Nosema maniereae* (Protozoa, Microspora) sur la biologie d'*Heliothis armigera* (Lepidoptère, Noctuidae)
- J. P. NENON et G. FABRES 605 Etude methodologique de l'efficacite parasitaire d'un Hymenoptera Encyrtidae neotropical *Epidinocarsis lopezi* introduit en Afrique pour lutter contre la cochenille du manioc *Phenacoccus manihoti*; bilan des travaux franco-congolais: 1982-1988

PART II

SYMPOSIUM ON INSECT PESTS AND SUSTAINABLE FOOD PRODUCTION

- K. F. NWANZE 615 *Plenary Lecture*
Sustainable food production in Africa: The challenge to national agricultural research systems (NARS)

ECOLOGY, BIOLOGY AND BIONOMICS

- E. N. MWANGI, R. M. NEWSON and G. P. KAAYA 629 Drop-off patterns for engorged adult females, nymphs and larvae of *Rhipicephalus appendiculatus*
- K. J. MBATA 635 Food plant preferences and spatial dispersion patterns of the armoured ground cricket, *Acanthopplus speiseri* Brancsik (Orthoptera: Tettigoniidae)
- S. B. SAGNIA, S. OUMAROU and A. HAROUNA 641 Preliminary results on the biology of the rice armyworm, *Mythimna loreyi* (Lepidoptera: Noctuidae)
- J. P. R. OCHIENG'-ODERO, F. O. ONYANGO, J. T. KILORI, M. D. O. BUNGU and E. O. AMBOGA 645 Insect rearing management as a prerequisite in the development of IPM for sustainable food production

K. V. SESHU REDDY

653 Insect pests of sorghum in Africa

INSECT PHYSIOLOGY

E. D. KOKWARO, L. H. OTIENO
and M. CHIMTAWI661 Salivary glands of the tsetse *Glossina pallidipes* Austen
infected with *Trypanosoma brucei* and virus particles:
Ultrastructural study

PEST MANAGEMENT, STRATEGIES AND PRACTICES

K. F. NWANZE

673 Components for the management of two insect pests of
pearl millet in Sahelian West Africa

J. ALLOTEY

679 Storage insect pests of cereal in small scale farming
community and their control

J. KAYITARE et L. NTEZURUBANZA

695 Evaluation de la toxicite et de l'effet repulsif de
certaines plantes du Rwanda contre les bruches du
haricot: *Acanthoscelides obiectus* Say et *Zabrotes*
subfasciatus Boheman

B. S. KILONZO

699 Larvicidal effects of neem, *Azadirachta indica*, on fleas
in Tanzania

S. M. K. J. LUGURU

703 Field observations on the residual effect of deltamethrin
against adult ticks, mainly *Rhipicephalus*
appendiculatus, during a high infestation period in the
Southern Province of Zambia

A. M. ALGHALI

707 The effect of plant spacings on cowpea, *Vigna*
unguiculata Walps, insect pests and yields in two sites
in Nigeria

A. DIAITE

713 La trypanosomiase animale Africaine au Senegal la lutte
anti-vectorielle

M. O. ODINDO

717 Potential of microorganisms for locust and grasshopper
control

Book Review

723

Instructions to Authors

i

Author Checklist

iv

SUBJECT INDEX

- A. acidula* 317
Abelmoschus esculentus 367–374
A. biplagae 317
Abrus precatorius 484
Abutilon sp. 323, 551
A. calamus 490
Acanthomia horrida 530, 590
Acanthopius speiseri 401–408, 409, 635–639
Acanthoscelides oblectus 695–697
Acanthospermum hispidum 405
Acerophagus advena 47
Acerophagus nr. *eric* 47
Aconus calamus 94, 487–491
Aceonurella sp. 349
Acorphy glaucopsis 718
Acholotermes chirotus 513
Acigona ignifusalis 654
Acremonium zonatum 356
 acridology 307
Acrocerops bifasciata 323
Acrotylus longipes incarnatus 718
A. curvipes 590
Acyrtosiphon pisum 65
Adelphocoris apicalis 321, 656
 adipokinetic hormones 279–286
Aedes aegypti 288–295
Aedes spp. 292
Aenasius advena 47
Aeneolamia albofasciata 65
Ae. vexans 189
A. fabae 353
A. fijiensis 344
 African armyworm 57–61, 530
 African Regional Pest Management
 Research and Development Network
 see also PESTNET
 Africa's NARS 616–617
Afrosteles distans 241
Afzelia africana 90
Agathis brusei 317, 321
Agathis sp. 40
Agave cantala 439–442
Agenionitellix deorum 205–208
Ageratum sp. 488
Agonoscelis haroldi 321
Agonoscelis pubescens 656
A. gossypii 530
A. granosus 202
 agricultural extension 579–583
Agrius convolvuli 530
 agromyzid beanflies 464–471
Agrostis setacea 205–208
Agrypon slaveolatum 13
Ahasverus advena 680–681
A. inconspicua 317
Aiolopus simulatrix 718
Alcornea cordifolia 714
Alestes sp. 361–365
 allele frequency 228–229
 allelopathic plant 356
 allelopathy 352
Allium cepa 94
Alphitobius diaperinus 680
Alphitobius laevigatus 571–578
Alstonia scholaris 353
Alternaria tenuis 351
Alternaria zinniae 351
A. luminosi 524
Amaranthus paniculatus 405
Amaranthus spp. 289–292, 235
Amascia sp. 324
Amblyomma hebraeum 629
Amblyomma variegatum 141–145, 704
Amblyseius aleyroides 316–338
Ambrosia pycnostachyum 205–208
 American bollworm 322
Ammora squamosa 94
Anablepia granulata 205–208
Anagyrus sp. 47, 316–338
An. arabiensis 375–378
Anastatus sp. 589–590
 anatomical parameters of 171–176
An. atroparvus 288–295
Ancylostoma duodenale 399
Andropogon gerardi 205–208
Anethum graveolens 94, 489
An. gambiae 292, 375–378
An. maculipennis var.
 sluius labbranchiae 290–292
Anopheles arabiensis 275–378
Anopheles maculipennis atroparvus
 290–291
Anopheles gambiae 531
Anopheles maculipennis var. *elutus*
 290, 292
Anopheles spp. 288–295
Anopheles stephensi 487
Anoplocnemis curvipes 321, 456, 460,
 589, 590
Anthermus granosus 201–208
 anthocorid bug 319
 antibiosis 319, 189–192
 antibiotic 352, 484
Anticarsia gemmatilis 66–67
 antifedant(s) 107
 action 94–95
 antixenosis 189–192, 449–454
Antrocephalus nr. *asubelongalus* 40
Antrocephalus sp. 35, 37
A. oblectus 90, 94, 95
Aonidomytilus albus 531
Apanteles angaletti 317
Apanteles biplagae 321
Apanteles dipaprosidis 317, 321
Apanteles flavipes 5, 19–26
Apanteles ruficornis 317
Apanteles sesamiae 647
Apanteles spp. 37, 317, 321, 589
Apanteles syleptiae 318, 324
A. parasiticus 178
Aphanogmus fijiensis 339, 342
Aphelopus sp. 316–338
 aphid control 117, 318
Aphidius ervi 277
Aphis craccivora 530, 557, 558, 561,
 708
Aphis fabae 350, 581–583
Aphis gossypii 349, 352
 control of 313–338, 547–552
 host of 385–389
Aphis spiraeicola 349
A. pisum 65
 application of insecticides 127–137
 techniques 111–125
 applied entomology 307
A. quadriannulatus 375
Arachnocampa luminosa 518
Aristolochia spp. 441
 armoured ground cricket 409
 armyworm 473–480, 641–643
Arphia conspersa 719
 artificial diet 209–214, 648
Ascaris lumbricoides 396, 399
Ascaris suum 399
Aschersonia spp. 64–65
A. scholaris 355
A. segax 318
Aspergillus flavus 178
Aspergillus spp. 364
A. stephensi 292
Astothrips bhatti 324
Atelosera spinulosa 321
A. tenuis 205–208, 354
Ateucholermes rastratus 513
Atherigona soccata 183–188, 653
A. tiplagae 317
A. tomentosicollis 530, 590
Autographa californicus 322
Azadirachta indica 94, 298
 for post-flowering plants 103–109
Babesia microti 383
Babesia spp. 379–384
Bacillus sp. 35, 37–41, 318
Bacillus thuringiensis 321–323, 326,
 530, 532, 675, 720
 for cereal pests 57–61
 isolation of 59–61
 baculovirus 326
 "bakars" 571–578
 banana weevil 531
Basidiobolus sp. 65
Bassia longifolia 94
Bassia talifolia 94
Bathypsectes curculianis 13
Batrachomorpha angustatus 349
B. bassiana 66, 67
Bauhinia rufescens 715
Beauveria bassiana 64–65, 369–370,
 372
Bemisia spp. 531
Bemisia tabaci 65, 319
 control of 547–552
 pest status of 315–338
 populations of 225–230
B. filipjevi 233
B. fusca 581–583
 BHC 530, 531

- biconical traps 380
Bidens pilosa 405–407
 bioassay 59
 biocontrol agents 35, 354
 biological control 6, 9–18, 311–312,
 326, 329, 339, 350, 353, 355–356,
 456, 529, 541–545, 588, 675, 688
 of cassava mealybug 45–49
 of cassava green mite 45–49
 of cowpea pests 585–592
 present and future prospects 3–8
 status of 347–359
 biological data 209–214
 methods of control 87–102
 biology of
 Goniozus sensorius 339–345
 Mythimna loreyi 641–643
 biometrical observations 419–422
 bionomics of 385–389
 biorational control 87–102
 biotechnology 63, 311
 biotypes 553, 559
Biplolaris tetramera 370–372
 black flies 531
Blaesoxipha rufipes 233
Blaesoxipha spp. 233
 blight 351
 blue disease 318–319
Bombyx mori 433–437
Boerhaavia 321
 boll development 322
 formation stages 321
 bollworms 313–338, 321
Boophilus decoloratus 141–145
Boophilus sp. 144
Boottitix punctatus 202, 205–208
 borer
 control with *Nosema* 52–53
 coffee berry control of 24–34
 management 54–55
 Bostrychidae infesting cereals 77–85
Bothriochloa insulata 405
B. pisorum 91
Bracon hebetor 676
Bracon sp. 37, 589
Brachaspis fulva 205–208
Brachaspis nivalis 205–208
Brachymeria feae 318
Brachymeria podagrica 233
Brachymeria sp. 317–318, 321, 324
Brachystegia spp. 141
Brachytripes sp. 324
Bracon kirpatricki 317
Brassica sp. 94
Braunsia sp. 37, 40, 589–590
 breeding for resistance 555
Brevipalpus phoenicis 350
B. rhodhani 383
 bruchid control 87–102
Bruchidius atrolineatus 88, 588
Bruchus brachialis 91
Bruchus pisorum 91
Bruchus rufimanus 88
Bruchus spp. 91
Bucculatrix parthenica 353
 buprofezin 391–394
Busseola fusca 51, 65, 187, 209, 530,
 654
 rearing of 645–651
Butyrospermum parkii 94
Cajanus cajan 385–389
Calidea dregii 656
Calidea sp. 321
Caliothrips funipennis 548
Caliothrips impurus 324
Callaophyllum inophyllum 94
Calliphora erythrocephala 518, 668
Callistemon lanceolatus 489
Callosobruchus chinensis 78, 487
Callosobruchus maculatus 88–89, 104,
 586, 588
Calocoris angustatus 183–188, 656
C. amada 490
Campanotus sp. 514
Campanotus sericeus 589, 590
Campoletis chloridae 369–372
Campoletis sonorensis 477
Camponotus rufoglaucus 37–41
Campophex haywardi 13
Campyloma nicolasi 321
Campylomma angestior 656
Canavallia sp. 215–223
C. apiculatus 65
Capsicum annuum 94
Capsicum frutescens 695–697
C. arachidis 249–261
 carbaryl 113
 carbon dioxide 288–295
Carebara vidua 511
C. argentea 299
Carpophilus dimidiatus 680
Carpophilus spp. 680
 carryover 353
Carthartus quadricollis 680, 682
 cassava green mite 43–49
 mealybug 43–49, 311, 605
Cassia nigricans 94
Cassia tora 353
Cassia uniflora 347, 352–353
Cataloipus cyruberiferus 718
Catantops sylvestris 202–208
Catantopsilus taeniolatus 202
Catopyla disorphaea 213
 cattle tick control 139–146
C. boninensis 318, 329
C. brunneus 530
C. camphora 488
C. canephora 154
C. caspici 354
C. cephalonica 682
C. chinensis 90
C. congrua 316–324
Cedria paradoxa 40
Cedrus deodara 490
C. elongata 456
Celosia 300
Cephus pygmaeus 13
C. excelsa 154, 156
C. flava 324
Chelonus blackburri 329
Ceiba maculatus 90–102
Centrosema pubescens 588
Cephalonica stephanoderis 34
Cercospora rodmanii 356
 cereals in storage 77–85
Ceroplastes sp. 349
C. erythrocephala 523
C. ferrugineus 680
 chaffy heads 674
Chalcodemus aeneus 587
Charops sp. 318
Chelonius insularis 473–480
Chelonius sp. 37
Chelonius versatilis 317
 chemical control 87, 348, 456, 541–
 545, 573, 576–577
 defense 104
 3, 52, 529, 530, 532, 667, 708–709
Chemopodium procerum 695
C. hypochlora 581–583
C. erythrocephala 523
C. harimansi 242–247
 vectors of 249–261
 chemosterilization 95
Chilo agamemnon 654
Chilo auriculus 19–26
Chilo diffusilineus 654
Chilo infuscatellus 19–26, 66
Chilo partellus 19–26, 311, 419–422,
 530, 557, 654
 control on sorghum 53–55
 evaluation for 53–55
 in maize 269–274
 management 51–55, 57–61
 rearing of 645–651
 resistant to 183–188
Chilo partellus resistance 147–151
Chilo sacchariphagus 19–26, 654
Chilo sp. 565
Chilo suppressalis 65
Chilo zacconius 599–604
Chilocorus angolensis 316–324
 chlorfenvinphos 320, 329, 704
 chlorinated hydrocarbon 190
 chlorpyrifos 320
Chorthippus brunneus 407
Chorthippus dorsatus 202, 205–208
Chorthippus montanus 202, 205–208
Chorthippus parallelus 205–208
Chromolaena odorata 324
Chrysocraon dispar 205–208
Chrysocoris stollii 349
Chrysopa boninensis 316–324
Chrysopa spp. 46–47, 317
Chrysopa vulgaris 316–338
C. ghaurii 238–247
 vectors of 237–247, 249–261
Cicadulina arachidis 238–247
Cicadulina leafhoppers 237–247
Cicadulina spp. 237–247, 249–261
Cicadulina triangula 238–247
Cicer arietinum 90, 367–374
Cidurcia awatocanda 318
 CILSS member countries 305–309
C. infuscatellus 65
Cinnamomum tamala 488
Clara magnifera 590
Clavigralla tomentosicollis 104, 588–
 590, 708
 cowpea antixenosis to 449–461
C. leucotreta 327, 581–583
C. liberica 154
C. longa 489

- Cissus adenocaulis* 321
C. mbila 581
 vectors of 249–261
C. nicolasi 656
 cob borer
 ecological data 215–223
 morphological and biological data 209–214
 taxonomic structures 238–247
 coccidia oocysts 395–400
 coccidiosis 399
Coccinella rufescens 316–324
Coccinia grandis 339
Coccobacillus acridiorum 720
Coccosterphus minutus 349
Coccus longulus 349
Codium sp. 482, 483
 codling moth 322
Coffea arabica 28, 95, 154–156
Coffea canephora 154–156
Coffea excelsa 154–156
Coffea liberica 154–156
Cola cordifolia 715
 collar rot 351
Collectotrichum capsici 350
Collectotrichum gloeosporioides 351
Collyria calcitrator 13
Colostriidium 37–41
Coniesta ignefusalis 673–678
Conidiobolus (Entomophthora) grylli 64–65
Conidiobolus spp. 65
 conservationists 6
Contarinia sorghicola 183–188
 control
 chemical measures 686
 contact insecticides 687
 of cotton insect 313–338
 cotton stainer 320
 methods 236
 of *Heliothis* 111–125
 of insect pests 19–26
 of mite pests 313–338
 of pests of sugar-cane 19–26
 physical measures 686
 practices 674–675
 strategies 231–236
 controlling of post-flowering pests 103–109
Corcyra cephalonica 441, 647, 680–681
Coriandrum sativum 489
C. orichalcociliellus 530, 654
Cordyla pinnata 715
Coryna spp. 581–583
Cosmophila flava 323, 315–338
Cosmopolites sordidus 531
Cotesia marginiventris 473–480
 cotton
 aphid 313–338
 insect 313–338
 in Zambia 111–125
 jassid 319
 pink bollworm 323
 whitefly 319
 cowpea insect pests 585–592, 707–711
 parameters of 171–176
 plant architecture 193–199
C. parazeae 250
C. pipiens quinquefasciatus 531
C. pudica 318, 324
C. puncticollis 530
C. quadricollis 680
Cremastus sp. 40
Creontiades pallidus 321, 656
 cricket 401
 crop hygiene 88
 protection training centre 305–309
 rotation 586, 587
 pattern 314
 cropping systems 177–182
Crotalaria juncea 178
Crotalaria spp. 588
Cryptocatanops haemorrhoidalis 718
Cryptolaemus montrouzieri 47
Cryptolestes ferrugineus 571–578, 681
Cryptolestes spp. 680
Cryptomenus bergi 178
Cryptophlebia leucotreta 209–214, 314–338, 317–338
Cryptophlebia (Argyroplote) leucotreta 322
Chrysopa malada boninensis 318
 crude oils 94
C. sacchariphagus 530
C. sericea 347
 see also *Cassia uniflora*
C. shadabi 456
C. similis 238–247
 vectors of 249–261
C. sordidus 531, 581–583
C. storeyi 241–247
C. subflava 656
C. subhyalinus 514
C. sudanensis 548
Ctenocephalides felis 593–597
Ctenocephalides spp. 596
C. testaceus 512
C. triangular 249–261
Cubitermes testaceus 511–514
 cultural aspects 87–102
 control 87–102, 456, 541–545, 549
 on cotton insects 313–338, 547–552
 on cowpea pests 585–592
 methods 456, 585–592
 practices 321, 323, 325, 531, 675, 710
 techniques 63
Culex 290
Culex nigripalpus 292
Culex spp. 288–295, 531
C. uniflora 353, 355–356
Curcubita maxima 385–389
Curcuma amada 487–491
Cuscuta reflexa 352
 cutaneous leishmaniasis 531
C. wahlbergi 316–324
Cx. thalassus 292
Cx. invidiosus 292
Cx. tarsalis 292
Cx. tigripes 292
Cx. triaeniorhynchus 292
Cydia ptychora 587–589, 710
Cydonia vicina 316–324
Cylas formicarius 530
Cylindrothorax westermanni 656
Cymbopogon flexuosus 488
Cynodon dactylon 475
 Cypermethrin 111–125, 127–137, 190, 323
Cyrus scariousus 485
Dactylis glomerata 205–208
 damage to crop plants 235
Datura stramonium 405
D. castanea 322, 325
 DDT 113, 530–531, 594
 dead heart 147–151, 183–188, 565, 647, 675
 deltamethrin 190, 327, 703–706
Dendrothrips fasciatus 324
Denntia tripetala 104
 density-damage relationships 270
Dentichasmias busseolae 647
Deracocoris aculatus 319
Deracocoris oculatus 316–338
Dermestes maculatus 361–365
 desert locust 71–75
Diabolocatanops axillaris 718
Diadegma cerophaga 13
 diamond-back moth 439–442
Diaperasticus erythrocephala 37–41
Diaphania indica 339–345
Diatraea saccharalis 5
 dictyophos 328
Digitaria sp. 405
 dimethoate 113, 320, 328, 530
Dinoderus dictinctus 77–85
Dinopsyllus lupus 593
Dionus sp. 47
Diopsis thoracica 710
Diparopsis 117, 137, 329
Diparopsis castanea 314–338
Diparopsis watersi 313–338, 321, 322
 dispersion patterns 409–417
D. melanoderes 320
Dociaustauous cruciger 205–208
Dolycoris indicus 349, 656
D. perditor 322
 drop-off patterns 629–633
Drosophila 377
Drosophila melanogaster 518, 522, 524
 dryinid parasitoid 319
D. superstitiosus 320
D. volkeri 656
D. watersi 322
Dysdercus 137
Dysdercus fasciatus 548
Dysdercus flavidus 320
Dysdercus koenigii 487
Dysdercus spp. 315–338
Dysdercus superstitiosus 650
Dysdercus sutirellus 453
Dysdercus volkeri 316–338
Earias 329
Earias biplaga 314–338
Earias insulana 314–338
Earias spp. 316–338
E. biplaga 321
Ecarsia lutea 316
Echium plantagineum
 ecological data 215–223
 economic control 104
 damage threshold 538

- impact of 139–146
 importance 673–674
 importance of 313–338
 injury level of 269–274
 losses 271
 threshold 318–319, 322, 325–326, 329
 ecosystems 511, 541–545
E. diversiciliatus 316
E. dolichi 319
 effect(s) of 591–597
 canopy structure 193–199
 crop phenology 53–54
 plant spacings on 707–711
 pod position 193–199
 solar radiation 53
 soybean hosts 189–192
 trap height 443–447
 effect(s) on bolls, buds and flowers 117
 yield 113
 mulches 297–303
 effectiveness of plant oils 77–85
 efficacy of entomopathogenic fungi 67
E. gayneri 656
Eichhornia crassipes 441
Eimeria arlongi 396–397
Eimeria oocysts 395–400
Elaeis guineensis 714
Eldana saccharina 51, 530, 654
 electrodyn
 sprayer(s) 111–125, 127–137, 328
 swath rows 127–137
 electrophoretic study of 275–278
 electrostatic sprayers 328
 elegant grasshopper 231–236
Eleusine indica 405
Elsholtzia polystachya 488
Ematheudes sp. 654
 embryonic development 95
Empoasca devastans 349, 352
Empoasca distinguenda 319
Empoasca dolichi 586
Empoasca lybica 547–552
Empoasca spp. 315–338
Empusa frensii 318
 emulsifier preparation 105
E. mundus 320
Encarsia lutea 320
Encarsia sp. 316, 320
Encotolophus sordidus 205–208
 encyrtid parasitoid 312
 endosulfan 328, 530
E. ninakohlyankimovae 396–397
Enterobacter cloacae 720
Entomophthora grylli 324, 719
 entomology for technicians 305–309
 entomopathogenic fungi 63–70
Entomophthora 64
Entomophthora planchoniana 65
 environmental manipulation 96
E. parva 396–397
Ephesia cauwella 680–682
Epiblema stenuana 350
Epidinocarsis lopezi 45–49, 312, 605–611
Epidinocarsis (=Apoanagyrus) lopezi 11
Epilachna verivestis 192
Epiricania melanoleuca 19
 epizootiology of 379–384
E. radicans 65
Eretmocerus mundus 316, 320
Eretmocerus sp. 320
E. rufocunealis 656
Erynia 64
Erynia neoaphidis 65
E. spathulacauda 202
E. sphaerosperma 65
E. strenuana 354
Eublemma brachygonia 656
Eucalyptus globulus 104
Eucalyptus sp. 353
Eucarcelia evolans 317, 322
Eucoptera anguliflava 201–208
Eucoptacra spathulacauda 201–208
Eupatorium japonica 488
Euphorbia balsamifera 714
Euphorbia heterophylla 405
Euphorus amcates 316
Euphorus sp. 316
Euplectrus laphygamae 317, 324
Eurystylus bellevoeyi 656
Eurytoma braconidis 318, 324
Eutetranychus africanus 325
 evaluation of
 bean cultivars 463–471
 bivoltine hybrids 433–437
 insect growth regulators 71–75
 juvenoid 71–75
Exitianus indicus 349
Exochomus nigromaculatus 316–324
Exochomus sp. nr. *sallaei* 47
 exposure to gases 78
Exserohilum rastratum 351
Extenterus amictorius 13
Extenterus vellicatus 13
Eyprepocnemis plorans 202–208

 factor analysis 423–431
 false codling moth 322
 farming systems 35–42, 139
 fecundity of 342,
 Bombyx mori 433–437
 Plutella xylostella 439–442
 Trioxys indicus 385–389
 Fenoxycarb 71–75
Ferrisia virgata 348, 349
Ferrisia virgata 320
 fertility of eggs 209–214
Ficus sycomorus 405
 Fish, dried damage to 361–365
 flea pest 593–597
 flowering stages 321
F. moniliforme 354–355
 food crisis 615–625
 plant preferences 635–639
 production 615–625
 foraging behaviour of
 Microtermes obesi 297–303
 Odontotermes spp. 297–303
Frankliniella schultzei 324
Frankliniella spp. 315–338
F. sjostedti 324
 fumigation 95
 fungal pathogens 160, 347
 for control of pests 63–70
 fungicides 543
Fusarium semitectum 351
Fusarium sp. 354
Fusarium moniliforme 352

G. aethiops 343
Gambroides javensis 13
 gamma radiation 95
G. arboreum 318
G. charon 590
G. claripennis 342–343
G. columbianus 343
G. cornutus 681
G. emigratus 343
 genetic control 87–102
 engineering 311
 expression of 147–151
 variation 225–230
 genitalia
 female 210–212
 male 210–211
 genotypes 104
Geocoris ruficeps 316
 geographical races 228–229
 germplasm accessions 456, 460
 collection of IITA 104
 germination test 68, 82–84
G. gallicola 343
G. fordhi 343–344
G. hirsutum 326
Gilpinia hercyniae 13
G. indicus 342, 343
G. japonicus 342–343
Glossina fuscipes fuscipes 662
Glossina longipalpis 713–715
Glossina morsitans morsitans 518, 523, 713–715
Glossina pallidipes 279–384, 661–669
Glossina palpalis gambiensis 713–715
Glossina spp. 531
 Glover mite 313–338
Glycine max 189–192
Glypta haesinator 13
G. marasmi 340–344
G. m. morsitans 524
G. montana 340
Gnatocerus spp. 680
G. naphantidis 343
Gomphrena globosa 352
Goniozus indicus 340
Goniozus naphantidis 340
Goniozus sensorius 339–345
Goniozus spp. 343
Gossypium arborium 326
Gossypium hirsutum 313–338
 gossypol content 323
G. pakmanus 343
G. palpalis 531
G. palpalis gambiensis 531, 715
G. raimondii 318
 grain legumes 530
 grainage parameters 433–437
 granaries types 96
Granulosis virusca 373
 grasshopper(s)
 control 717–722
 feeding 201–208
 niche breadth 201–208
 greater grain borer 311
 groundnut intercropping 177–182

- growth index 189–192
 growth regulator 439–442
Gryon charon 589
Gryon gnidus 589–590
G. sensorius 340, 342
G. tachinoides 531
G. thurberi 326

Haemaphysalis bispinosa 629
Haemaphysalis leporispalustris 629
Haemoproteus columbae 668
 Hardy-Weinberg 375, 377
H. armigera 66, 322–327, 581–583
Harungana madagascariensis 202–208
Hayetiola destructor 539
H. bicinctus 581–583
H. citriflorus 64, 66, 67
H. cymbaria 636–639
H. dissoluta 636–639
 head bugs resistant to 183–188
Hecalus arcuatus 349
Helicoverpa (Heliothis) armigera 322–323
 population dynamics 367–374
Heliocheilus albipunctella 673–678
Heliothis 127–137, 329
Heliothis armigera 209, 656, 350, 599–604,
 control of 313–338, 547–552
Heliothis zea 559
Heliothis control 111–125
Heliothis spp. 66
Helminthosporium sp. 356
Helminthosporium tritici-repentis 354
Heliopeltis schowedeni 126–338
Hepalia machaeralis 40
 herbicides 543
Hercinothrips bicinctus 531
H. eruditus 353
Heteracris guineensis 202–208
 heterosis 147–151
 heterobeltiosis 433–437
Heteropsylla cubana 4
 heterosis 433–437
H. filipendula 636–639
H. fleischeri 322
Hibiscus spp. 323
Hibiscus cannabinus 405, 551
Hibiscus esculentus 405, 551
Hibiscus sabdariffa 551
Hilda partruelis 530
Hirsutus thompsonii 67
Hirsutiella sp. 65
Hirsutiella thaupsonii 64
Hishimonus phycitis 349, 352, 355
H. nyassae 636–639
 host plant resistance 160
 host preference in 287–295
 seeking of 287–295
 range 444
 housefly 487–491
H. madagascariensis 202–208
H. nitidulus 581–583
H. rufa 636–639
H. senegalensis 316–324
H. suaveolens 355
H. truncatum 142
 human resource constraint 619
H. virescens 559

Hyalomma spp. 141
Hyalomma excavatum 629
Hyalomma truncatum 704
 hybrids 433–437
 Hymenoptera parasitic 9–18
Hyparaspis notata 47
Hyparrhenia 141, 416
Hyparrhenia bracheata 405, 636–639
Hyparrhenia cymbaria 405
Hyparrhenia dissoluta 405
Hyparrhenia filipendula 405
Hyparrhenid nyassae 401–408, 635–639
Hyparrhenia rufa 405, 407
Hyp. diandra 205–208
Hypera postica 13
Hyperaspis notata 46
Hyperaspis pumiia 316–324
Hyperaspis sp. 47
Hypothenemus eruditus 348, 350
Hypothenemus hampei 27–34
Hypoestes verticillaris 203–208
Hyptis spicigera 94
Hyptis suaveolens 353

Icerya australis 324
Icerya nigroareolata 153
Icerya pattersoni 4, 153–157
Icerya seychellarum 348, 349
 Ichneumonidae African 9–18
Illicium verum 487–491
 immunity 383
 inert substances 90
 infestation levels 313
 injury threshold 373
 insect bioecology 674
 growth regulators 71–75
 incidence of 361–365
 in Nigeria 707–711
 of sorghum 653–657
 of cereal 679–693
 pest management 708
 pests 263
 rearing management 645–651
 resistance 183–188
 insecticidal
 properties of 487–491
 action 94–95
 control 233–236
 insecticides
 application 111–125, 127–137
 on cotton 111–125, 127–137
 locally produced 104
I. sulphurea 153
 integrated
 biological control programme 348
 control 324
 management programmes 3
 pest control 547–552
 pest management (see IPM) 87, 96, 108, 325, 443–447, 535–539, 541–545, 553–564
 intercropping 63, 586
 effects of 177–182
 interspecific competition 473–480
 intrinsic rate of 385–389, 419–422
Ipomoea 300, 301
Ipomoea fistulosa 297–303
 IPM 5–6, 35, 269–274, 535–545
 components 646–651
 IPM development 645–651
 packages 645–651
Isaniris decorsei 324
Ischiodon scutellaris 316–324
Isoberlinia spp. 141
I. sulphurea 153
 irradiation processes 95
 irrigation 63
I. verum 490
 Iwao's index 409, 411
 regression 412–413, 414–415
Ixodes persulcatus 629
Ixodes ricinus 629

Jacobiella (Empoasca) facialis 319
Jacobiella (Empoasca) lybica 319
 jassid control 117
Jatropha curcas 94
 JH 72–75, 286
 JHA 94–95
Julbernadia spp. 141
 juvenile hormones 78
 juvenile hormone analogue 94–102
 juvenoid insect growth regulators 71–75

 kairomone 289, 454
Kalodiplosis sp. nov. 47
 karyotype 375, 377
 knapsack sprayers 111–125, 327
Kotschyia sp. 405

Lablab niger 550
 ladybird beetle 4
Lagenaria vulgaris 385–389
 lambda cyhalothrin 111–125
Lantana 225–230
Lantana camara 353, 405
Larrea tridendata 202, 205–208
Laspeyresia nigricana 13
Latheticus oryzae 680
 leafhoppers 249–261, 319
 leaf roller 323
 leafworm 323
Leucopis (xenoleucopis) cilifemur 47
 legume pod borer
 damage by 193–199
 infestation by 193–199
 resistance to 171–176
 leaf spot 351
 legislative regulations 547–552
Leofa sp. 349
Leptocentrus taurus 348–349
Lespedeza cuneata 205–208
Leucania psyllid 4
Leucania separata 642
Leucas martinicensis 405, 407
Leucas stricta 405
 levels of resistance 159–169
 life table(s) 385–389, 423–431
 life table data 35
 light traps 445–446
 lipoprotein interconversions 279–286
Lippia multiflora 201–208
Listronotus setosipennis 355
L. multiflora 202–208
 locust
 control 717–722
 flight 279–286
Locusta migratoria 280–282, 308

- Locusta migratoria migratorioides* 717–722
- Locusta pardalina* 718
- Loudeia simplex*
- L. rivularis* 656
- L. taurus* 353
- Luffa cylindrica* 385–389
- Luperomorpha vittata* 348, 349, 353
- Lycopersicon esculentum* 367–374
- Lygaeus pandurus* 656
- Lygus lineoris* 559
- Lygus* spp. 316–338
- Lygus (Taylorigus) vosseleri* 321
- Lygus vosseleri* 315–338
- Machaeridia bilineata* 205–208
- Macrotermes estherae* 301
- Macrostelas fascifrons* 518
- Macrotermes subhyalinus* 513
- Macrotermes thoracalis* 548
- Mahanarva posticata* 66–67
- maize
- germplasm 159–169
 - streak virus 249–261
- Malamoeba locustae* 718
- Maliarpha separattella* 710
- Maliarpha* sp. 565
- malpighian tubules 517–525
- Mamestra brassicae* 321
- management
- of *Chilo partellus* 51–55
 - of pest(s) 77–85, 673–693
- Mangifera indica* 714
- Manihot esculenta* 11, 43–49
- M. anisopliae* 66, 667
- Margarodes* sp. 320
- marine algae 481–485
- Maruca* 456
- Maruca testulalis* 557, 586, 589
- control of 103–109
 - damage by 193–199
 - infestation by 193–199
 - natural enemies 35–42
 - natural populations of 423–431
 - resistance to 171–176
- mass
- formulations 63
 - production 63
- Massospora* 64
- mathematical model 391–394
- Mayetiola destructor* 554
- M. bivittatus* 718
- M. brassicae* 322–323
- M. dawsoni* 718
- M. differentialis* 718
- M. domestica* 523–524
- mealybug 4, 312
- medical vectors 531
- Megalurothrips sjostedii* 104, 175, 588–590, 708
- Meganoptera foetens* 511
- Melaleuca leucadendron* 489
- Melampsora lini* 556
- Melanaphis sacchori* 65
- Melanoplus bivittatus* 202, 205–208, 719
- Melanoplus differentialis* 718
- Melanoplus fermurrubrum* 202, 205–208
- Mendel's laws 554
- Menida distantis* 656
- mephosfolan 531
- Mermis nigrescens* 719
- Mermis* sp. 316
- Metarrhizium flavoviridae* var. *minus* 233
- Metarrhizium* sp. 317
- Meteoridae testacea* 324
- Meteorus testaceus* 317
- Meteosus laphygmarum* 317, 321, 323
- methamidophos 320
- Metharizium flavoviride* 67
- methods of bruchid control 87–102
- Mettriona circumdata* 349
- Mentha* spp. 94
- Mesoleius tenthredinis* 13
- Messor barbarus* 653
- Metarrhizium anisopliae* 64, 66
- Meteoridae testacea* 318
- Mettriona circumdata* 349
- Mettesia* 37–41
- M. flavoviride* 66
- M. fragrans* 490
- microsporidia
- management with 51–55
 - for field application 54–55
- Microtermes obesi* 178, 299–301
- behaviour of 297–303
- Microtermes* spp. 177–181
- Microtermes thoracalis* 653
- midge resistant to 183–188
- migration activity of 252–261
- mildew 354
- militaristic approach 311
- Mimacrotoma cingulata* 47
- Mimosa pigra* 715
- mirid bugs 321
- Mitragina inermis* 714
- Microbracon brevicornis* 323
- microorganic pathogens 35
- Microplitis croceipes* 477
- Microplitis manilae* 473–480
- Mirperus jaeulus* 656
- Miscanthus sinensis* 205–208
- mite
- cotton 325
 - pests 318–338
 - red spider 325
 - yellow tea 325
- mitochondria 517–525
- mixed cropping 586
- modelling 264
- Molinia coerulea* 205–208
- monocrotophos 328
- monoculture 587
- Monolepta signata* 349
- Monomorium* sp. 589
- Mononychellus* spp. 65
- Mononychellus tanajoa* 43–49, 530, 649
- morphogenic damage 71
- morphological data 209–214
- mortality rate(s) 35, 108
- mosquito attractant 288–289
- mosquitoes, host-seeking of 287–295
- moth migration 322
- M. parkardii* 719
- Mucuna pruriens* 202–208
- mulches on foraging 297–303
- multiple cropping 88
- insect resistance 183–188
- Musca domestica* 40, 396, 399, 487–491, 518
- Musca* spp. 395–400
- Mussidia nigripenella* 323, 681
- biological data 209–214
 - ecological data 215–223
 - morphological data 209–214
- M. vicina* 399
- mycoplasma like organism 352
- mycoplasmal disease 350–351
- mycotoxins 350, 355–356
- Mycterotherips acaciae* 324
- Mylabris* spp. 581–585, 656
- Myristica fragrans* 487–491
- Myrmelatettix maculatus* 205–208
- Mysotra dilecta* 324
- Mythinna loreyi* 641–643
- Nagasta* sp. 589–590
- national agricultural research 615–625
- natural
- control of arthropod populations 63–70
 - control of entomopathogenic fungi 64–70
 - enemies 96, 339, 353–354, 368, 539
 - enemies conservation of 22
 - enemies for control 19–26
 - enemies of pest 43–39
 - enemies in pest management 5–6
 - enemies of pod borer 35–42
 - increase 419–422
 - mortality 232–233
 - populations 423–431
 - products 490
 - resistance 88
- nature of resistance 556
- Necrobes rufipes* 363
- neem cake 297–303
- neem oil 103–109
- emulsifiable concentrate 103–109
 - extraction 104
- nematode 35
- Nematospora* sp. 320
- Neochetina eichhorniae* 356
- Neodiprion swaini* 13
- Neorthacris simulans* 349
- Neoseilus teke* 645–651
- Neozygites fumosca* 64
- Neozygites* sp. 65
- Nephotettix cincticeps* 275
- Nephotettix modulatus* 319
- Nephotettix nigropictus* 275
- Netelia* sp. 317
- Nezara viridula* 321, 656
- N. fumosa* 65
- Nicandra physaloides* 401–408, 636–639
- Nilaparvata lugens* 65–67, 275, 391–394, 559
- Nipaeococcus corymbatus* 349
- Nola analis* 656
- Nomadacris septemfasciata* 717, 718
- Nomuraea rileyi* 66, 67
- nonadditive gene 147
- nonpreference 676
- Nosema locustae* 717–722
- Nosema manierae* 599–604
- Nosema maruca* 718

- Nosema* sp. 35, 37–41, 51–55
for *Chilo partellus* 53–55
evaluation of 53–55
- Nosema pyrausta* 52
- Nosema* spores 52–53
- nuclear polyhedrosis virus 373
- Oberea* sp. 349
- Ocimum basilicum* 488
- Ocimum gratissimum* 487–491
- Ocimum sinus* 695–697
- Ocyptamus* sp. 47
- Odontomachus haematoda* 512
- Odontotermes obesus* 299
- Odontotermes* spp. 299, 177–181, 302
behaviour of 297–303
- Oedaleus nigeriensis* 718
- Oedaleus senegalensis* 308, 718
- Oedipoda coerulescens* 205–208
- Oencyrtus* spp. 589
- O. haematoda* 514
- O. gratissimum* 488, 490
- Oidium parthenii* 351
- Oidium* sp. 351
- O. insignis* 353
- O. kilimandscharicum* 488
- Olesicampe benefactor* 13
- olfactory chemosensilla 288–295
- olfaction
in host-seeking 287–295
role of 287–295
- Oligonychus gossypii* 44
- Olla quinquengrum* 47
- O. mercator* 681
- Omethoate* 320
- Onchocerca volvulus* 531
- onchocerciasis 531
- on-farm storage 159–169
- Oothea* beetles 707–711
- Oothea benningseni* 586
- Oothea mutabilis* 589, 590
- O. parthenii* 354
- Operophtera brumata* 13
- O. phaseoli* 581–583
- Ophiomyia phaseoli* 463–471
- Ophiomyia spencerella* 463–471
- Ophiotermes ugandensis* 513
- Opinus* sp. 589
- Opisina arenosella* 5
- Orchochla brachycnemis* 205–208
- organophosphate 190
- Orobancha aegyptiaca* 352
- Orosinus albicinctus* 349
- Orosinus albicinctus* 352
- Orosinus cellulosus* 319
- Orseola oryzivora* 566
- Orthezia insignis* 348, 349
- Orthochlha brachycnemis* 202
- Orthogalumna terebrantis* 356
- Oryctes rhinoceros* 67
- Oryza sativa* 679–693
- Oryzaephilus surinamensis* 680
- O. sanctum* 488
- O. senegalensis* 718
- Ostrinia nubilalis* 52, 150, 214
- O. surinamensis* 683, 685
- O. ugandensis* 514
- ovicide 94
- oviposition 232, 235, 342
- oviposition preference 450, 454, 469
- oviposition sites 90
- O. wallonensis* 299, 301
- Oxya velox* 205–208
- Oxycaenus hyalinipennis* 321
- Oxyrhaechis tarandus* 349
- Oxythenantera abyssinica* 715
- Oryzaephilus surinamensis* 681
- Pachnola cordata* 656
- Paecilomyces fumosoroseus* 64
- palm oil 77–85
- Palorus* spp. 680
- Palorus subdepressus* 571–578
- Paltothyrsus* spp. 512, 514
- Paltothyrsus tarsus* 514
- P. americana* 524
- Paprides australis* 205–208
- Paprides nitidus* 205–208
- Parachrysocharis javensis* 19–26
- Paragus borbonicus* 316–324
- Parahieroglyphus bilineatus* 205–208
- Paramixia saturalis* 656
- Parapleurus alliaceus* 205–208
- Parasaissetia nigra* 348, 349
- Parasierola* sp. 340
- parasitic hymenoptera 9–18
- parasitoid(s) 35, 233, 235, 322
- Parkia africana* 90
- Parthenium hysterophorus* 347–359
- Patanga succinta* 65
- pathogen 350–351
- Paulina pinnata* 715
- Payrocephala gossypii* 320
- Pectinophora* (*Platyhendra*) *gossypiella*
314–338, 323, 548
- Pediculus humanus* 596
- Pediobius amorocoeus* 324
- Pediobius furvus* 647
- Penicillium* spp. 364
- Pennisetum glaucum* 673–678
- Pennisetum tiphoides* 88, 507–509
- percentage weight loss 81–82, 84
- Peribaea* sp. 323, 369–371, 372
- Perilampus mawrus* 233
- Periplaneta americana* 581
- Periplaneta* sp. 104
- Permanent Interstate Committee for
Drought Control in the Sahel
see also CILSS
- permethrin on 593–597
- pest
carryover 322
control 311, 536, 553, 579–583,
585–592, 679–693
control techniques 3
ecology 313–338
management 313, 321, 368,
529–533, 553–564, 586–587, 544
management natural enemies in 5–6
management with *Bacillus*
thuringiensis 57–61
resistance 553–564
species 311
status of 231–236
- pesticide 328–329
abuse 581
regulation 7
use 581
- pesticides 313, 323, 327, 535–539,
490–491
- PESTNET 68, 238, 312
- pests
of cassava 43–49
of cereals 679–693
cereal management 57–61
control with fungal pathogens 63–70
of cultivated crops 311
of pearl millet 673–678
of stored products 311
in tropics 63–70
- P. farinosus* 64
- P. fumosoroseus* 67
- P. gossypiella* 323, 327
- Phanerotoma* sp. 40, 589, 590
- Phaseolus funatus* 215–223
- Phaseolus lunatus* 88, 405
- Phaseolus mungo* 90
- Phaseolus vulgaris* 89–102, 197, 463–
471, 648
- Phaulacridium vittatum* 720
- Pheidole megacephala* 322
- Pheidole* spp. 589, 590
- Phenacoccus manihoti* 4, 11, 43–49,
64–65, 311–531, 605–611
- Phaeolophus basizomus* 13
- pheromone traps 443–447
- pheromones 95
- Philodocus temerarius* 316
- Phoetaliotes nebrascensis* 719
- Phonoctonus fasciatus* 589–590
- Phonoctonus lutescens* 317
- Phonoctonus nigrofasciatus* 316, 321
- Phthorimaea operculella* 13, 442
- P. hysterophorus* 350–351
- Phyllocoptruta oleivora* 67
- phyllody 351–352
- physical control 87–102
- planting density 63
- Piliostigma thornningii* 405
- P. interrupta* 656
- Piper guineense* 94
- Piper nigrum* 94
- P. irritans* 596
- Planococcus kenya* 320
- Planococcus* sp. 348–349, 353
- plant
architecture 193–199
associations 401–408
breeding 553–564
essential oils 487–491
genetics 553–564
growth regulations 327
oils effectiveness of 77–85
- planthopper 391–394, 441
- P. latus* 325
- P. lilacinus* 67
- Plodia interpunctella* 680–681
- Plutella maculipennis* 13
- Plutella xylostella* 6, 65, 439–442
- P. marginata* 656
- PMDISS 312
- P. nigra* 348
- P. nigrofasciatus* 317
- Podagrica puncticollis* 548
- Podagrica* sp. 324
- Podagrixena pallida* 324

- pod borer 456
 control of 103–109
 natural enemies 35–42
 natural populations of 423–431
 pod-sucking bug 449–454
 pod wall barrier 91
Podostemon plectranthoides 488
 polyhedrosis virus 323
Polyphagotarsonemus
 (*Hemitarsonemus*) *latus* 313–338
Polyspilota aeryginosa 589–590
Pongamia glabra 94
Poppilia japonica 415
 population
 dynamics of 367–374
 fluctuations 218–223
 growth of 391–394
 interchange 401–408
 post-flowering pests 103–109
 post-harvest loss 683
 post-independence period 617
 powdery mildew 351
P. pallida 548
 precision test 264
 prediction 266
 predators 35
 preference test 190–191,
 450–454
Pristiphora erichsonii 13
Proceras sacchariphagus 13
 production economics 139
Prorops nasula
 for control 27–34
 use of 27–34
Prospaltella sp. 316
Prostephanus hominivorax 311
Protenomus anoplocmedis 589
Pseudarthria sp. 405
Pseudolynchia canariensis 668
Psalydolyta fusca 543, 656
Pseudococcus sp. 350
Pseudomonas solanacearum 350
Pseudoperichaeta insidiosa 40
Pseudoperichaeta laeis 589
Pseudophonocortus sp. 590
Pseudoprotacta burmeisteri 656
Pseudosphinogonatus savignyi 718
Psidium guajava 104
P. simplex 656
P. subdepressus 680
P. theresae 656
P. tripola 656
Ptochus ovulum 350
P. truncatus 596, 680, 681, 682, 683
Puccinia abrupta 354
Pulex irritans 593–597
 purified oils 94
Pyrgomorpha cognata 718
Pyricularia oryzae 543
Pyrilla perpusilla 19–26
Pyroderces homizopa 656
Pyroderces simplex 323
 quanalphos 328
 quarantine 7
Radolia cardinalis 4
Raghava albipunctella 543, 673
Raphia sp. 715
Rastrococcus invadens 312
R. carmelita 589, 590
R. dominica 681–682
 rearing methodology 649
 recurrent selection methods 147–151
 red bollworm 321
Regas sp. 317
 reinfestation 78, 83
 repellent(s) 104, 107
 action 94–95
 properties of 487–491
 residual effect of 703–706
 resistance 439, 455–461, 463–471
Chilo partellus 147–151
 cultivars 45, 326, 554
 index 458
 of pests 328
 to *Sitophilus zeamais* 159–169
 sorghum 557
 varieties 90, 95, 529, 539, 550, 687
Rhabdophea munda 205–208
Rhabdosia mellisoid 488
Rhabdosia sobrina 656
Rhapalosiphum maidis 65
Rhinocoris albopilosus 559, 590
Rhipicephalus appendiculatus 141–
 145, 629–633, 703–706
Rhipicephalus evertsi 704
Rhipicephalus spp. 141–145
Rhinocoris bicolor 589–590
Rhizoctonia solani 351
Rhizopertha dominica 571–578, 680–
 682
 suppression of 77–85
Rhizopus spp. 364
Ricinus communis 561
Riptortus dentipes 460, 589, 590, 708
R. maidis 581–583
 root crops 530
 root rot fungus 351
Rottboelia 416
Rottboelia exaltata 401–408, 475,
 635–639
R. risbeci 656
Ryacionia frustrana 13
Saba senegalensis 715
Sacrophaga dispersa 317
Saissetia coffeae 348–349
 salivary glands of 661–669
 sampling 252–261, 411
S. calamistis 654
 scale, cushion 4
 scanning electron microscope
 237–247
 scarification 178
Schistocerca americana, 71–75
Schistocerca gregaria 205–208, 308,
 548, 719
 control of 717–720
Schistocerca pellenis 720
Scirpophaga nivella 13
Sclerotium rolfsii 351
S. coarctata 67
Scolothrips hartwigi 324
Scotinophora coarctata 67
 screening assays 59
 of *P. nasula* 31
Scymnus ornatus 316–324
S. damnosum 531
S. derogata 324
S. dysderci 317
 seed damage 449–461
 seed treatment 319
Selebria mesozonella 656
Selenops radiatus 589–590
 semi loopers 323
 semi-synthetic diet 648
Sericothrips occipitalis 586
 serotypes 321
Serratia marcescens 720
Sesamia albivena 654
Sesamia botanophaga 209–214, 654
Sesamia calamistis 51, 209–214, 530,
 566, 654
Sesamia cretica 654
Sesamia nonagrioides 654
Sesamia penniseti 654
Sesamia poephaga 654
Sesamia sp. 565
Sesamum orientale 94
Sesamum orientale 352
Setaria sphacelata 407, 636–639
Setaria verticillata 405
 sex attractant 443–447
 sex pheromones 322–323, 326
 sex-ratio 209–214
S. exigua 66, 317–338
S. frugiperda 66–67
S. galilaenus 361–365
S. granarius 104
S. hamata 355
 shoot-fly resistant to 183–188
Shorea robusta 94
Sida cordifolia 319
Sida rhombifolia 405
 silkworm 433–437
Sitophilus zeamais 78, 104, 682
 resistance to 159–169
Sitophilus granarius 681
Sitophilus oryzae 571–578, 682
Sitophilus spp. 680
Sitotroga cerealella 507–509, 656,
 674, 681–684
 control of 571–578
S. littoralis 66, 323
 small scale farming 679–693
Smicronyx lutulentus 353
 socio-economic aspects 87, 104
Sogatella furcifera 275–278
Sorghum spp. 679–693
Sorghum vulgare 405, 635–639
S. oryzae 104
 spatial dispersion patterns 635–639
 spatial genetic variation 225–230
Spartina alterniflora 205–208
 species identification 238–247
Sphenoptera sp. 324
Sphenostylis stenocarpa 215–223
Spherophthera gossypii 548
 spider mites 350
 toxicity of 481–485
Spilarcia casignia 189–192
Spilosoma obliqua 348, 350
Spilostethus spp. 656

- spinning disc sprayer(s) 113, 127
 spiny bollworms 321
Sogatodes oryzicola 554
Solanum dubium 551
Solanum nigrum 405
Sonchus oleraceus 405
Sorghum bicolor 475, 653
 sorghum genotypes 183–188
Sorghum spp. 679–693
 soybean
 hairy caterpillar 189–192
 hosts development on 189–192
Spodoptera exempta 719
 management 57–61
Spodoptera frugiperda 67, 107, 473–480
Spodoptera littoralis 315–338, 323, 547–552, 719
 management 57–61
Spodoptera litura 66, 443–447
Spodoptera spp. 317–338
Spodromantis lineola 589–590
Sporobolus pyramidalis 405
 spotted stem borer 147–151, 311
S. rhombifolia 319
S. sudanensis 316–324
 stable fly 517–525
Stathmopoda auriferella 656
 stem
 borer(s) 350, 354, 419–422, 529
 borer cereal management 51–55
 borer in maize 269–274
 borer resistant to 171, 183–188
 borer spotted 147
 tunneling 674
Stenocrobylus festinus 202–208
Stobaera concinna 353
Stomoxys calcitrans 517–525
Stomoxys sp. 65, 290
 storage
 facilities 160
 hermetic 89
 insect pests 307, 679–693
 pits 571–578
 techniques 96
 stored grain insects 571–578
 storing unthreshed pulses 89
S. oryzae 680–681
Striga hermonthica 543
Stylosanthes fruticosa 353
Stylosanthes lamata 352
 sugar-cane insect pests 19–26
 sunnhemp intercropping 177–182
 survival curves 35
 sustainable production 615–625
Sylepta derogata 315–338
Symphorobius sp. 47
 synergistic response 289
 synthetic pyrethroids 190
 syrphids 318
Systates sp. 581–583
 systems approach 139–146
S. zeamais 680–681, 683
- literature 12
 structures 237–247
Tayloriylgus vosseleri 656
 Taylor's Power Law 409, 411–415
T. brasiliensis 317, 323
T. brucei 379–384
T. brucei brucei 713
T. brucei gambiense
T. castaneum 680, 682–683
T. confusum 680
T. congolense 379–384
 technicians at the CILSS 305–309
Telingana campbelli 349
Temelucha difficilis 474
Temnoschoila nigropagiata 65
 temperature regulation control 89
 temporal genetic variation 225–230
Tenebrio molitor 680
Tenebroides mauritanicus 680
 termite(s) 511–514
 behaviour of 297–303
 damage 177–182, 300
 incidence 177–182
Tetradentia viparia 695–697
Tetranychus cinnabarinus 325, 350
Tetranychus indeni 350
Tetranychus macfarlanei 350
Tetranychus neocaledonicus 350
Tetranychus spp. 315–338
Tetranychus telarius 481
Tetranychus urticae 325, 481–485
Tetrapleura tetraptera 103–109
Tetrastichus atriclavus 317–318
Tetrastichus sesamiae 37, 647
Tetrastichus sp. 589
Thaia sp. 349
Theileria spp. 379–384
Thelairosoa palpsoum 588–589
Thelohania sp. 668
Therioaphis maculata 560
 thiodicard 329
Thripocetus sp. 316
 thrips 531
Thrips tabaci 324
Thuja occidentalis 488
Thymus serpyllum 487–490
 tickborne disease 139, 140
 tick control 139–146, 629–633
 tick-sensitized animals 631
 timing of harvesting 88
Tirathaba complexa 13
Tithonia rotundifolia 405, 636–639
T. neocaledonicus 325
T. niloticus 361–365
 tolerant varieties 539
 toxicity of 481–485
 toxins 104
Trachyspermum ammi 489
 training
 in entomology 305–309
 manuals 308
 trap-cropping 586, 588
 traps 443–447
 treatment with synthetic insecticides 78
Triaspis thoracica 91
Tribolium castaneum 571–578, 680, 681–682
Tribolium spp. 680
Tribulus terrestris 442
Trichodesma sp. 352
Trichogramma 326
Trichogramma chilonis 19–26
Trichogramma evanescens 317, 323
Trichogramma mwanzai 645–651
Trichogramma sp. 316, 318
Trichospilus diatraea 317
Trichoplusia ni 66, 192
Trichophisia orichalcea 350
Tridacnoides tardus 277
Tridax procumbens 94
Trioxys indicus 385–389
Trimerotropis saxatilis 202, 205–208
Tristachya soperia 405
Triticum spp. 679–693
Trogoderma granarium 680
Trombicula spp. 596
 tropical grasshoppers 201–208
Tryachypogon spicatus 405
Trypanosoma brucei 661–669
Trypanosoma congolense
Trypanosoma spp. 379–384
Trypanosoma vivax 379–384, 713
 trypanosomiasis 379–384, 531
Tryporyza incertulas 263–267
T. serpyllum 490
T. sudanensis 316
T. telarius 325
Typhlodromus fallacis 481
Typhlodromus medianicus 316
- Ultra-low volume sprayers 111–125
 ULV spinning disc sprayers 128
 ULV sprayers 327–328
Uscana semifumipennis 91
U. senex 91
Utethesia pulchella 178
- varietal resistance 313–338, 676
 vector-borne diseases 311
 vector(s) 249–261
 management 529–533
 varietal control 87–102
 vedalia beetle 3–8
 vegetable oil(s) 77–85, 94
 vegetative stages 321
 veins characteristics of 153–157
Venturia palmaris 13
Vernonia cinerea 405
Vernonia pertusifolia 405
Verticillium lecanii 64, 66, 719
Vestula lineaticeps 589–590
 virus particles 661–669
 visceral leishmaniasis 531
Vicia faba 88, 94
Vigna radiata 91
Vigna unguiculata 36, 103–109, 455–461, 586, 707–711
 canopy structure 193–199
 parameters of 171–176
Vitiellaria paradoxa 215–223
- weed control 588
 weevil 159–169
 whitefly 117, 225–230
 whitehead 565

wilt 351
witches' broom 352

Xanthodes graelsii 323
Xanthogramma pfeifferi 316
Xanthomonas malvacearum 320
Xanthomonas parthenicola 351
Xanthopimpla sp. 318
Xenopsylla braziliensis 593
Xenopsylla cheopis 593, 596
X. parthenicola 354

yellow tea mite 313–338
yield loss 313–338

Zabrotus subfasciatus 88–89, 102,
695–697
Zanthoxylum alatum 489
Z. bicolorata 353–354, 356
Zea mays 405, 475
 Acanthopplus speiseri on 635–639
 Helicoverpa armigera on 367–374
 storage pests of 679–693

Zea elegans 324
Zingiber elatum 489
Zonocerus 235
 Zonocerus elegans 66, 231–236
 Zonocerus spp. 233
 Zonocerus variegatus 44, 65, 104, 308,
 324, 531, 718
Zoophthora 64–65
 Z. phalloides 65
 Z. subfasciatus 94
 Z. variegatus 108, 231–236
Zygomma bicolorata 347–359

AUTHOR INDEX

- Agrawal, B. L. 183
 Agwaro, P. O. 35
 Ahmed, M. A. 547
 Alghali, A. M. 707
 Allotey, J. 679
 Amboga, E. O. 645
 Amer, S. A. A. 481
 Ashley, T. R. 473
 Awoyemi, M. D. 361

 Bacye, B. 565
 Baliddawa, C. W. 579
 Beenackers, A. M. Th. 279
 Bellotti, A. 225
 Bhatt, N. 385
 Bowers, W. S. 71
 Brownbridge, M. 57
 Bungu, M. D. O. 645

 Chiang, H. S. 455
 Chintawi, M. 661
 Chizyuka, H. G. B. 139
 Cwikla, P. S. 237

 Dabrowski, Z. T. 237, 249
 Dakouo, D. 565
 David, B. V. 339
 Devaiah, M. C. 433
 Devaraj Urs, C. 439
 Diaite, A. 713
 Dimetry, N. Z. 481
 Dipeolu, O. O. 395

 El Tigani, M. El A. 547
 Ezueh, M. I. 585

 Fabres, G. 605

 Gahukar, R. 313
 Gerling, D. 225
 Gezahegn, T. 375
 Gold, C. S. 177, 297
 Gopalan, M. 391
 Govindan, R. 433
 Giga, D. P. 159
 Greathead, D. J. 3
 Gupta, V. K. 9

 Harouna, A. 641
 Harris, K. M. 311

 Jackai, L. E. N. 103, 171, 193, 455
 Javaid, I. 111, 127

 Kaaya, G. P. 629
 Kailasam, C. 391
 Kayitare, J. 695
 Kayumbo, H. Y. 463
 Khan, V. 395
 Kilonzo, B. S. 593, 699
 Kilori, J. T. 645
 Kinuthia, M. W. 153
 Kokwaro, E. D. 661
 Kongoro, J. 517
 Kumar, R. 77, 249

 Lavigne, R. J. 571
 Luguru, S. M. K. J. 703
 Lulu, M. 375

 Makanjuola, W. A. 171, 193
 Mainoya, J. R. 517
 Maniania, N. K. 63
 Mazarura, U. M. 159
 Mbata, K. J. 409, 635
 Mohyuddin, A. I. 19
 Morales, F. 225
 Moyal, P. 209, 215
 Mueke, J. M. 463
 Murphy, S. T. 27
 Mwangi, E. N. 629

 Nacro, S. 565
 Narayanaswamy, K. C. 433
 Nenon, J. P. 605
 Nesbitt, S. T. 379
 Neupane, F. P. 189
 Newson, R. M. 629
 Nderitu, J. H. 463
 Nigatu, W. 375
 Njau, B. C. 379
 Nolt, B. 225
 Ntezurubanza, L. 695
 Nwanze, K. F. 183, 615
 Nwilene, F. 249
 Nyambo, B. T. 231

 Ochieng'-Odero, J. P. R. 645
 Odebiyi, J. 449
 Odhiambo, T. R. 1, 507, 517
 Odindo, M. O. 51, 29, 717
 Oghiakhe, S. 171, 193
 Okeyo-Owuor, J. B. 35, 423
 Okonronkwo, N. O. 77
 Okwakol, M. J. N. 511
 Olatunde, G. O. 455
 Oloo, G. W. 35, 423

 Onyango, F. O. 645
 Oosterwijk, G. P. M. 139
 Ortego, F. 71
 Otieno, L. H. 379, 661
 Oumarou, S. 641
 Oyediran, I. O. 103

 Pathak, R. S. 147, 553
 Pegram, R. G. 139
 Peter, C. 339
 Philippe, L. G. 201
 Pimbert, M. P. 297
 Prasad Reddy, G. V. 439
 Pushpalatha, N. A. 347

 Rajapakse, R. H. S. 473
 Rajendram, G. F. 275
 Ranga Rao, D. V. 443
 Rangi, D. K. 27
 Reda, A. S. 481
 Reddy, Y. V. R. 183
 Ri-Zhi, L. 263

 Sagnia, S. B. 305, 641
 Salin, K. P. 391
 Seshu Reddy, K. V. 269, 653
 Sharma, H. C. 183
 Singh, A. K. 487
 Singh, D. 487
 Singh, R. 367, 385
 Singh, S. P. 419
 Srikanth, J. 349
 Sum, K. O. S. 269

 Takken, W. 287
 Taneja, S. L. 183
 Tilahun, D. 375
 Toguebaye, B. S. 599
 Tran, M. 209, 215
 Tripathi, S. R. 367
 Tukahirwa, E. M. 535

 Umeh, E-D. N. 43

 Van Huis, A. 87
 Visweswara Gowda, B. L. 433

 Waddill, van H. 473
 Wanjala, F. M. E. 153
 Wightman, J. A. 177, 297, 443
 Wool, D. 225

 Zethner, O. 541
 Zheng-Xiang, Y. 263

INSECT SCIENCE AND ITS APPLICATION

The International Journal of Tropical Insect Science

AIMS AND SCOPE

Insect Science and its Application deals comprehensively with all aspects of scientific research targeted on tropical insects (and related arthropods), and the application of new discoveries and innovations to such diverse fields as pest and vector management and use of insects for human welfare.

A distinctive feature of the journal is its multi- and inter-disciplinary nature, which transcends the traditional boundaries of entomology.

Its second feature is its concentration on the recording and reviewing of the progress of insect science in the tropical and subtropical regions of the world. Thus, without excluding any area of the wide-ranging field encompassed by insect science, the journal will accept manuscripts in environmental physiology, the regulation of development and reproduction, population modelling, chemical ecology, natural products chemistry, plant resistance, host-insect relations, behaviour of tropical pest species, epidemiology of tropical diseases, vector biology, pest and vector management research, insect pathology, entomo-meteorology, insects in relation to farming systems, forest entomology, social insects and arthropods, and the use and farming of insects. It is the intention of the Editors to have the manuscripts published rapidly, consistent with the needs of quality control.

In addition to articles of original research, the journal also publishes mini-review articles, announcements and reports of meetings, book reviews, new patents, obituaries of prominent scientists and software survey. Each regular issue of the journal usually contains a short review article on a critical or rapidly developing area of tropical insect science; normally, the Editors will have invited an author to contribute such a mini-review.

Insect Science and its Application is an international journal, pursuing its own independent policy through its editorial staff and the Editorial Advisory Board, which is

internationally appointed. Its sponsors, the ICIPE and the African Association of Insect Scientists (AAIS), do not intervene in editorial matters.

INSTRUCTIONS TO AUTHORS

For indications of the design of papers, the author should consult a recent issue of the journal. The following paragraphs are provided as guidelines for the preparation and submission of manuscripts.

1. All papers for publication in *Insect Science and its Application* should be submitted directly to Professor Thomas R. Odhiambo, Editor-in-Chief, *Insect Science and its Application*, ICIPE Science Press, P.O. Box 72913, Nairobi, Kenya. Papers should be mailed in a strong envelope, preferably linen, as they will otherwise arrive in a damaged condition.
2. Papers should be typed in English or French, and be original contributions in the field of tropical insect science. Each paper should have a summary in the form of an abstract not exceeding 200 words in both English and French.
3. Submission of a paper implies that it has not been published previously, that it is not under consideration for publication elsewhere, and that if accepted for *Insect Science and its Application* the authors will transfer copyright (by signing the form) to the ICIPE Science Press of the International Centre of Insect Physiology and Ecology. Manuscripts and illustrations become the property of the journal.
4. There is no page charge for papers accepted for publication.
5. Manuscripts and illustrations must be submitted in triplicate (original and two copies) to ensure efficient refereeing. If a computer is used for the preparation of the manuscript, the diskette should also be submitted (the author should state the hardware and software used and ensure that it is virus-free). In the case of multiple authorship, the authors should indicate who is to receive correspondence and the correct address for the mailing of proofs.
6. Manuscripts must be typewritten with double spacing (including the reference list), with wide margins, and on one side of the paper only. Authors are requested to keep their

communications as concise as possible. Footnotes should be avoided, and italics should not be used for emphasis. A running head of not more than 30 letters should be supplied.

7. The manuscript should contain the following features:

Title: This should be followed by the author's name and full address.

Abstract: Containing a brief summary of the contents, and conclusions of the paper. It should not be longer than 200 words, and should be supplied in both English and French.

Key Words: A short list of key words (arranged in alphabetical order) should be provided for rapid scanning of the contents of the paper and use in compiling the Index to the completed volume of the journal.

Introduction: Should contain a brief survey of the relevant literature and the reasons for doing the work.

Materials and Methods: Sufficient information should be given to permit repetition of the experimental work by other workers. The technical description of methods should be given only when such methods are new or modified.

Results: Should be presented concisely. Only in exceptional cases will it be permissible to present the same set of results in both a table and a figure. The results section should not be used for discussion.

Discussion: Should be separate from the Results section, and should deal with the significance of the results and their relationship to the object of the work.

Comparison with relevant published work should be made, and conclusions drawn.

Acknowledgements: Only pertinent acknowledgements should be cited.

Reference: The Harvard system will be followed. References should be detailed in the following order: Authors' names and initials, date of publication (in brackets), the title of the article, the name of the journal as abbreviated in the *World List of Scientific Periodicals* (4th edn. 1964), the volume, and the first and last pages of the article, e.g.

Delobel A. G. L. (1983) Influence of temperature and host plant condition on preimaginal development and survival in sorghum shootfly *Atherigona soccata*. *Insect Sci. Applic.* 4, 327-335.

For books, the author's names, date of publication, title, edition, page reference,

publisher's name and the place of publication should be given, e.g.

Zurflueh R. C. (1976) Phenylethers as insect growth regulators: laboratory and field experiments. In *Juvenile Hormones* (Edited by Gilbert L. L.), pp. 61-74. Plenum Press, New York.

Within the text, references should be given as Unnithan (1981) or (Unnithan, 1981). When a citation includes more than two authors, e.g. Ochieng', Onyango and Bungu, the paper should be referred to in the text as Ochieng' et al., provided that this is not ambiguous. If there are only two authors, they should be referred to as Ochieng' and Onyango. If papers by the same authors in the same year are cited, they should be distinguished by letters (a,b,c, etc.).

Reference to papers "in press" must mean that the article has been accepted for publication and should be given as follows:

Darlington J. P. E. C. (1988) Primary reproductives and royal cells of the termite, *Macrotermes michaelsoni*. *Insect Sci. Applic.* (In press).

References to "personal communication" and unpublished work are permitted in the text only.

8. Illustrations. All necessary original illustrations should accompany the manuscript, but should not be inserted in the text. All photographs, graphs and diagrams should be numbered consecutively in arabic numerals in the order in which they are referred to in the text.

Black and white glossy photographs or positive prints (not slides or X-ray negatives) should be sent, unmounted wherever possible, and should be restricted to the minimum necessary.

Charts, graphs or diagrams should be drawn boldly in black ink on good quality white paper or card. Lettering to appear on the illustrations should be given in full and should be of sufficient size to withstand considerable reduction, where necessary. Alternatively, lettering may be shown on a rough overlay and will be inserted by the printer.

Illustrations should not be larger than 15 x 24 cm, or they may be damaged in the post. On the back of each illustration should be indicated the author's name, the figure number (in arabic numerals), and the top of the illustration, where this is not clear.

The following symbols should be used on

line drawings as they are readily available to the printer: ▲ △ ▼ ▽ ■ □ × + ● ○.

Legends to figures should be typed on a separate sheet, not under the figures, and should give sufficient data to make the illustration comprehensible without reference to the text.

Magnification of photographs and illustrations should be indicated by an appropriate scale bar and not in the legend. This practice obviates the complications which arise through change of size in reproduction.

9. Tables should be constructed so as to be intelligible without reference to the text, numbered in arabic numerals, and typed on separate sheets. Every table should be provided with an explanatory caption, and each column should carry an appropriate heading.

Units of measurement should always be indicated clearly. All tables and figures must be referred to in the text in numerical order.

10. Only standard abbreviations should be used. Where specialized abbreviations are

used, the name should be given initially in full with the abbreviation indicated in parentheses.

11. The common and Latin names should be given for all species used in the investigation, although taxonomic affiliation and authority should not necessarily be provided in the title.

12. Page proofs will be sent to the authors by the publisher. They will already have been proof read by the proof reader for the typesetter's errors; but it is the author's responsibility to ensure that the proof contains no errors. Correction to proofs should be restricted to typesetter's errors or editorial queries only. Other than these, substantial alterations will be charged to the author. Proofs should be returned to the ICIPE Science Press within 48 hr of receipt for final preparation for printing. Failure to meet this deadline may result in the inadvertent inclusion of errors in the published article. The publisher will not be held responsible for printed errors, if the author overlooks them.

